



# Discrete Event Systems

## Solution to Exercise Sheet 6

### 1 Gloriabar

- a) You might be tempted to model this situation by a queue with a bounded number of states, because the maximal number of persons in the line is bounded by 540. However, the situation can also be modeled by an infinite M/M/1 queue without losing too much accuracy; the parameter  $\rho$  will not be too large, such that the probability to reach the state in which 540 persons are standing in the queue at once is extremely small anyway. The modeling by an infinite M/M/1 queue conveniently allows us to apply Little's Law and therefore, we can use the formulae for the response and waiting time:

Our arrival rate  $\lambda$  and service rate  $\mu$  are given by

$$\lambda = \frac{540}{90 \cdot 60} = \frac{1}{10}, \quad \mu = \frac{1}{9}$$

(persons per second). Thus  $\rho = \frac{\lambda}{\mu} = \frac{9}{10}$ . By Theorem 5.19, the expected number of persons in the M/M/1 system is


$$N = \frac{\rho}{1 - \rho} = \frac{\lambda}{\mu - \lambda} = 9.$$

By applying Little's Law, we learn that the expected time until the student has paid for her food is  $T = \frac{N}{\lambda} = 90$  seconds. The expected waiting time is  $W = T - \frac{1}{\mu} = \frac{\rho}{\mu - \lambda} = 81$  seconds.

- b) We use the formula for the expected number of jobs in the queue and obtain queue length of  $N_Q = \frac{\rho^2}{1 - \rho} = 8.1$ .
- c) We require that  $T = \frac{N}{\lambda} = \frac{1}{\mu - \lambda} = \frac{90}{2}$ , where  $\lambda = 0.1$ . Thus,  $\mu = \frac{11}{90}$ , i.e., instead of 9 secs, the service time should be  $\frac{90}{11} \approx 8.2$  secs.

### 2 Beachvolleyball

- a) We know that the minimum of  $i$  independent and exponentially distributed (with parameter  $\lambda$ ) random variables is an exponentially distributed random variable with parameter  $i\lambda$ . Thus, we have the following birth-death-process:



b) Let  $\pi_i$  be the probability of state  $i$  in the equilibrium. We know that

$$\pi_i = \pi_0 \cdot \prod_{j=0}^{i-1} \frac{\lambda_j}{\mu_{j+1}}$$

and thus

$$\pi_i = \pi_0 \cdot \frac{\lambda_0 \cdot \lambda_1 \cdots \lambda_{i-1}}{\mu_1 \cdot \mu_2 \cdots \mu_i}.$$

Applying this formula to our process yields

$$\pi_i = \pi_0 \cdot \frac{n(n-1) \cdots (n-i+1) \cdot \lambda^i}{1 \cdot 2 \cdots i \cdot \mu^i} = \pi_0 \cdot \binom{n}{i} \cdot \rho^i \quad (1)$$

where  $\rho := \frac{\lambda}{\mu}$ . We know that the sum of all probabilities equals 1, so we have

$$\sum_{i=0}^n \pi_i = \pi_0 \sum_{i=0}^n \binom{n}{i} \rho^i = 1$$

Using the given formula for the binomial series

$$\sum_{i=0}^n \binom{n}{i} x^i = (1+x)^n$$

we obtain

$$\pi_0(1+\rho)^n = 1.$$

Finally, we obtain

$$\pi_i = \frac{\binom{n}{i} \rho^i}{(1+\rho)^n}.$$

c) (i) It is  $\rho = 3/9 = 1/3$ . We calculate the probability that there are less than two fit players:

$$\begin{aligned} \pi_0 + \pi_1 &= \frac{1}{(1+\rho)^n} \cdot \left(1 + \binom{n}{1} \cdot \rho^1\right) \\ &= \left(\frac{3}{4}\right)^5 \cdot \left(1 + \frac{5}{3}\right) \\ &= \frac{3^5}{2^{10}} \cdot \frac{8}{3} \\ &= \frac{3^4}{2^7} \approx 0.63 \end{aligned}$$

Thus, the DISCO team cannot participate in the tournament with probability 0.63.

(ii) Now,  $\rho = 2/4 = 0.5$ . Again, we calculate  $\pi_0 + \pi_1$ :

$$\begin{aligned} \pi_0 + \pi_1 &= \frac{1}{(1+\rho)^n} \cdot \left(1 + \binom{n}{1} \cdot \rho^1\right) \\ &= \frac{1}{1.5^5} \cdot (1 + 0.5 \cdot 5) \\ &= \frac{2^5 \cdot 3.5}{3^5} \approx 0.46 \end{aligned}$$

Hence, the probability that the DISCO team cannot participate is 0.46!

(iii) In general, if  $\rho \geq 1$ , an M/M/1 queue might grow infinitely and therefore doesn't have a stationary distribution. This cannot happen in this birth-and-death process, though, because there is only a bounded number of states. Hence, the process has a stationary distribution even for  $\rho \geq 1$ .

### 3 Theory of Ice Cream Vending

The situation can be described by a classic M/M/2 system. There is an equilibrium iff

$$\rho = \lambda/(2\mu) < 1 .$$

For the stationary distribution, it holds that

$$\begin{aligned}\pi_0 &= \frac{1}{\left(\sum_{k=0}^{m-1} \frac{(\rho m)^k}{k!}\right) + \frac{(\rho m)^m}{m!(1-\rho)}} \\ &= \frac{1}{\frac{(2\rho)^0}{0!} + \frac{(2\rho)^1}{1!} + \frac{(2\rho)^2}{2!(1-\rho)}} \\ &= \frac{1}{1 + 2\rho + \frac{4\rho^2}{2(1-\rho)}} \\ &= \frac{1}{1 + 2\rho + \frac{4\rho^2}{2(1-\rho)}} \\ &= \frac{1}{\frac{2(1-\rho) + 4\rho(1-\rho) + 4\rho^2}{2(1-\rho)}} \\ &= \frac{2(1-\rho)}{2 - 2\rho + 4\rho - 4\rho^2 + 4\rho^2} \\ &= \frac{2(1-\rho)}{2 + 2\rho} \\ &= \frac{1-\rho}{1+\rho} .\end{aligned}$$